

## PEAK EXPIRATORY FLOW RATES

**Suggested reference peak expiratory flow (PEF) values for children:**


Height (cm)	PEF		PEF	
	Caucasian		African*	
	Male	Female	Male	Female
100	127	142	120	126
101	131	145	124	130
102	135	149	128	133
103	138	152	131	137
104	142	156	135	140
105	146	159	139	144
106	150	163	143	148
107	154	166	147	151
108	158	170	151	155
109	162	174	155	159
110	166	178	159	163
111	170	182	163	167
112	175	185	168	171
113	179	189	172	175
114	184	193	176	179
115	188	197	181	184
116	193	202	186	188
117	197	206	190	192
118	202	210	195	197
119	207	214	200	201
120	212	218	205	206
121	217	223	210	210
122	222	227	215	215
123	227	232	220	220
124	232	236	226	225
125	237	241	231	230
126	243	245	236	235
127	248	250	242	240
128	254	255	248	245
129	259	259	253	250
130	265	264	259	255
131	271	269	265	260
132	276	274	271	266
133	282	279	277	271
134	288	284	283	277
135	294	289	289	282
136	300	294	295	288
137	307	299	302	293
138	313	304	308	299
139	319	309	315	305
140	326	315	322	311
141	332	320	328	317

Height (cm)	PEF		PEF	
	Caucasian		African*	
	Male	Female	Male	Female
142	339	325	335	323
143	345	331	342	329
144	352	336	349	335
145	359	342	356	342
146	366	348	363	348
147	373	353	371	354
148	380	354	378	361
149	387	365	386	368
150	395	371	392	374
151	402	377	401	381
152	410	382	409	388
153	417	388	417	395
154	425	394	425	402
155	433	401	433	409
156	440	409	441	416
157	448	413	442	423
158	456	419	458	430
159	464	426	466	437
160	473	432	475	445
161	481	438	484	452
162	489	445	492	460
163	498	451	501	468
164	506	458	510	475
165	515	465	520	483
166	524	471	529	491
167	533	478	538	499
168	542	485	548	507
169	551	492	557	515
170	560	499	567	523
171	569	506	577	532
172	578	513	587	540
173	588	520	597	548
174	597	527	607	557
175	607	534	617	566
176	617	541	627	574
177	626	549	638	583
178	636	556	648	592
179	646	563	659	601
180	657	571	670	610

\*Based on African American data.

For optimal control, 80% of the predicted peak flow is required.

### Peak expiratory flow in normal adult subjects



*Adapted with permission from Nunn AJ Gregg I, Br Med J 1989;298;1068-70 and Clement Clarke International.*

**CALCULATING % PREDICTED PEAK FLOW RATE**

- Take the best of 3 of the patient's observed peak flow rates (l/min):  
e.g. 200, 180, 190 performed – so take 200.
- Find the patient's sex, age and height predicted value from the nomogram.  
e.g. 440 l/min for a woman of age 25 years and height 167 cm
- Divide patient's observed peak flow rate over their predicted peak flow rate: e.g.  $200/440 = 0.45$
- Multiply by 100: e.g.  $0.45 \times 100 = 45\%$

So, in this example, the patient's observed peak flow rate is 45% of their predicted.

**CALCULATING BRONCHODILATOR RESPONSIVENESS USING PEAK FLOW IN ADULTS**

Perform peak flow testing and select the best of the 3 values to use as the pre-bronchodilator peak flow.

- Administer salbutamol 400 µg using a metered dose inhaler and spacer without a mask.
- Wait 15 minutes before repeating peak flow
- Repeat peak flow testing to obtain a post-bronchodilator peak flow.
- Subtract the pre-bronchodilator reading from the post-bronchodilator reading.
- Divide the difference by the pre-bronchodilator reading.
- Multiply by 100.

For example, a patient with readings that improve from 300 to 400, has reversibility of 33%. Measurements that improve by >20% strongly suggest a diagnosis of asthma. (See Sections 16.1: Asthma, acute and 16.2: Asthma, chronic persistent).

**CALCULATING PEAK FLOW VARIABILITY IN CHILDREN AND ADULTS**

- Perform peak flow measurements 4 times per day spread over the course of the day.
- Subtract the lowest reading of each day from the highest reading.
- Calculate the mean/average reading by adding all 4 readings from that day and dividing total by 4.
- Calculate PEF variability:

$$\text{PEF variability} = \frac{(\text{Highest PEF} - \text{Lowest PEF})}{\text{Mean PEF}} \times 100.$$

Determine this value on each day over two weeks, and average the results. Excessive diurnal PEF variability defined as >10% in adults and >12% in children strongly supports a diagnosis of asthma.

**ASTHMA CONTROL TEST™**

This is a validated measure of clinical asthma control that can be completed by the patient (after initial instruction) at each visit to the clinic prior to consultation. A value of  $\geq 19$  suggests adequate asthma control.

Online version of the test is accessible at: <https://www.asthmacontroltest.com/>

*Reference: Nathan RA, Sorkness CA, Kosinski M, Schatz M, Li JT, Marcus P, Murray JJ, Pendergraft TB. Development of the asthma control test: a survey for assessing asthma control. J Allergy Clin Immunol. 2004 Jan;113(1):59-65. <http://www.ncbi.nlm.nih.gov/pubmed/14713908>*